



International Journal of Gerontology

journal homepage: <http://www.sgecm.org.tw/ijge/>



CME for 18.4

- Which of the following conditions or comorbidities is most likely to predict emergency department visits in home healthcare patients?
(A) Depression
(B) Diabetes
(C) Having urinary catheters
(D) Hypertension
(E) Osteoarthritis
- Which of the following factors is likely to be associated with all-cause mortality in Japanese community-dwelling individuals?
(A) Body mass index
(B) Exercise habits
(C) Gender
(D) Handgrip strength
(E) All of the above
- Which of the following statements regarding cholangiocarcinoma is true?
(A) Cholangiocarcinoma is the second most common primary liver cancer after hepatocellular carcinoma.
(B) Cholangiocarcinomas can be categorized into three subtypes: intrahepatic, perihilar, and distal.
(C) Biliary drainage is an intervention for obstructive jaundice to relieve biliary tract obstruction and associated discomfort.
(D) Biliary drainage can be performed by percutaneous transhepatic cholangial drainage or endoscopic retrograde biliary drainage.
(E) All of the above.
- To quantitatively define sarcopenia, which of the following methods can be used to measure the skeletal muscle mass index?
(A) Computed tomography (CT)
(B) Magnetic resonance imaging (MRI)
(C) Dual-energy X-ray absorptiometry (DEXA)
(D) Bioelectrical impedance analysis (BIA)
(E) All of the above
- Which of the following factors can affect the burden of care for family caregivers of the elderly with dementia?
(A) Social support
(B) Degree of dementia
(C) Family monthly income
(D) Educational level
(E) All of the above

Issue date: October 1st, 2024

Expiration date: January 31st, 2025

Volume 18 Issue 3 Answers:

- (B)
- (B)
- (C)
- (D)
- (C)