

International Journal of Gerontology

journal homepage: http://www.sgecm.org.tw/ijge/



CME for 19.1

- 1. According to the report in "Does the Timing of Depression Onset in Individuals with 2-Year Survival CAD Affect Mortality? A Nationwide Retrospective Study", which was NOT a significant risk factor for all-cause hospitalization among CAD individuals with depression?
 - (A) Gender
 - (B) Age
 - (C) Income
 - (D) Comorbidities
 - (E) Depressive disorder (after one month group)
- 2. Which of the following statements in "The Effects of MST on Nursing Home Residents with Sleep Disturbance" is WRONG?
 - (A) Nursing home residents often experience sleep disorders
 - (B) MST is Magnetic Stimulation Therapy
 - (C) Cognitive function can assess by Mini-Mental State Examination Scale
 - (D) MST was effective for improving rapid eye movement sleep time
 - (E) MST was not an effective method to improve the objective sleep quality of nursing home residents with sleep distur-
- 3. According to the article "Lithium Toxicity Related Parkinsonism in an Older Woman: Case Report and Literature Review", which of the following statement is WRONG?
 - (A) Lithium is commonly used to treat affective disorders
 - (B) The most severe manifestations of lithium toxicity involve the central nervous system and kidneys, and some of these effects can be permanent
 - (C) Younger adults eliminate lithium more slowly and are at higher risk of lithium toxicity than are older adults
 - (D) The toxic effects of lithium mainly affect the kidneys, liver, heart, and glands but are usually mild and reversible
 - (E) Lithium serum level of > 1.0 mmol/L may cause neurological symptoms, and prolonged lithium intoxication can lead to permanent brain damage

- 4. What was **NOT** the limitation mentioned in the study of "Preliminary Results of Femoral Neck System in Treating Femoral Neck Fractures in Elderly Population"?
 - (A) It was an observational and retrospective study design
 - (B) The study lacked a control group for comparison
 - (C) Most of variables do not exhibit statistically significant differ-
 - (D) Clinical and radiological outcomes were documented with a small patient cohort
 - (E) FNS was released only in 2018, and it has been gradually used in our institute since 2022
- 5. According to the article "Pneumonia in Elderly Patients with Nasogastric Tubes: Factors to Consider" which was NOT a significant risk factor for community-acquired pneumonia (CAP) in elderly patients with nasogastric tubes?
 - (A) Age
 - (B) Female gender
 - (C) Impaired consciousness
 - (D) Excessive poly-pharmacy
 - (E) A signed DNR

Issue date: January 1st, 2025 Expiration date: April 30th, 2025

Volume 18 Issue 4 Answers:

- 1. (C)
- 2. (E)
- 3. (E)
- 4. (E)
- 5. (E)